

**The Alcohol Use Disorders Identification Test Consumption screening tool  
(AUDIT-C)**

Patient Name: \_\_\_\_\_

Date: \_\_\_\_\_

Alcohol consumption may affect your health and interfere with certain medications. Therefore, it is important that we assess your alcohol consumption by asking the following questions:

1. How often do you have a drink containing alcohol?

☐ Never (0)

☐ 2 - 3 times a week (3)

☐ Monthly or less (1)

☐ 4 or more times a week (4)

☐ 2 - 4 times a month (2)

2. How many standard drinks containing alcohol do you have on a typical day? A standard drink equals to 1.5 oz of liquor (i.e., whisky cocktail), a 12 oz beer or 5 oz of wine.

☐ 1 or 2 (0)

☐ 7, 8 or 9 (3)

☐ 3 to 4 (1)

☐ 10 or more (4)

☐ 5 to 6 (2)

3. How often do you have six or more drinks on one occasion? A standard drink equals to 1.5 oz of liquor (i.e., whisky cocktail), a 12 oz beer or 5 oz of wine

☐ Never (0)

☐ Weekly (3)

☐ Less than monthly (1)

☐ Daily or almost daily (4)

☐ Monthly (2)

**Scoring and Interpretation:** The number enclosed by parenthesis represents the score assigned to each response. Add the values the response in each question to obtain the total score. Refer to the following to determine alcohol misuse:

- Men: score of 4 points or more
- Women: score of 3 points or more

**Total Score:** \_\_\_\_\_